

[more quarks]

MQ-1

[About 1965]

Leptons

$\nu_e$   $\nu_\mu$   
 $e^-$   $\mu^-$

Quarks

u  
d s

Glashow proposed a new "charmed" quark c [wouldn't it be charming?]  
w/ a property called charm C

[Nov 1974] New meson discovered  $\psi$  /  $m \sim 3000 \text{ MeV}$  [ $\rightarrow$  see plot.]

called  $\begin{cases} \psi & \text{[SLAC, Richter]} \\ J & \text{[BNL, Ting]} \end{cases}$

$\downarrow$  [PPB, toward end of mesons, p. 116 in 2018 edition]  
 $J/\psi = \text{charmonium} = c\bar{c}$   $m \sim 3100 \text{ MeV}$

$\Rightarrow m_c \sim 1500 \text{ MeV}$

Other charmed mesons soon discovered

$D^+ = c\bar{d}$   
 $D^0 = c\bar{u}$  }  $m \sim 1900 \text{ MeV}$

$D_s^+ = c\bar{s}$   $m \sim 2000 \text{ MeV}$

Charmed baryons

Group these  $\left[ \begin{array}{l} \Sigma_c^{++} = cuu \\ \Lambda_c^+, \Sigma_c^+ = cud \\ \Sigma_c^0 = cdd \\ \Xi_{cc}^{++} = ccu \\ \Xi_{cc}^+ = ccu \\ \Xi_{cc}^0 = ccd \text{ and } \Xi_{ccc}^{++} = ccc \end{array} \right.$  [analogous to  $\Sigma \sim \begin{smallmatrix} suu \\ sud \\ sdd \end{smallmatrix}$ ]





Fig. 10.23. Total hadron production cross section in  $e^+e^-$  collisions near 3.1 GeV and the  $J/\psi$  peak. [From A. M. Boyarski et al., *Phys. Rev. Lett.* **34**, 1357 (1975).]


[ New quantum number  $\Rightarrow$  another dimension to weight diagram ]



Use hypercharge Y instead of S

$$Y = S + A - \frac{1}{3}C$$

|   | $I_3$          | A             | S  | C | Y              |
|---|----------------|---------------|----|---|----------------|
| u | $\frac{1}{2}$  | $\frac{1}{3}$ | 0  | 0 | $\frac{1}{3}$  |
| d | $-\frac{1}{2}$ | $\frac{1}{3}$ | 0  | 0 | $\frac{1}{3}$  |
| c | 0              | $\frac{1}{3}$ | 0  | 1 | 0              |
| s | 0              | $\frac{1}{3}$ | -1 | 0 | $-\frac{2}{3}$ |



Quarks belong to  $\underline{4}$  of  $SU(4)$

Mesons belong to  $\underline{4} \otimes \overline{\underline{4}} = \underline{15} \oplus \underline{1}$

$\uparrow$   
cuboctahedron

baryons belong to  $\underline{4} \otimes \underline{4} \otimes \underline{4} = \underline{20} \oplus \underline{20}_S \oplus \underline{20}_A \oplus \overline{\underline{4}}$

$\uparrow$  pyramid       $\underbrace{\hspace{2cm}}$  truncated pyramids       $\uparrow$  inverted tetrahedron




Figure 14.4: SU(4) multiplets of baryons made of  $u$ ,  $d$ ,  $s$ , and  $c$  quarks. (a) The 20-plet with an SU(3) octet. (b) The 20-plet with an SU(3) decuplet.




Figure 14.1: SU(4) weight diagram showing the 16-plets for the pseudoscalar (a) and vector mesons (b) made of the  $u$ ,  $d$ ,  $s$  and  $c$  quarks as a function of isospin  $I$ , charm  $C$  and hypercharge  $Y = S + B - \frac{C}{3}$ . The nonets of light mesons occupy the central planes to which the  $c\bar{c}$  states have been added.

$\Sigma_c^+, \Lambda_c^+$  ~~uuc~~ cud  
 $\Sigma_c^{++}$  cuu  
 $\Sigma_c^0$  cdd  
 $\Xi_{cc}^{++}$  ccu  
 $\Xi_{cc}^+$  ccd  
 $\Omega_{ccc}^{++}$  ccc

$D^0$   $c\bar{u}$   
 $D^+$   $c\bar{d}$   
 $D_s^+$   $c\bar{s}$  (1976)

Third generation

|         |           |            |             |
|---------|-----------|------------|-------------|
| $\nu_e$ | $\nu_\mu$ | $\nu_\tau$ | (disc 1999) |
| $e^-$   | $\mu^-$   | $\tau^-$   | (disc 1975) |
| $u$     | $c$       | $t$        | (disc 1994) |
| $d$     | $s$       | $b$        | (disc 1977) |

top/bottom

truth/beauty

(1977)  $\Upsilon = \text{upsilon} = \text{bottomium } b\bar{b}$   $m \sim 10 \text{ GeV} \Rightarrow m_b \sim 5 \text{ GeV}$

(1983) B-mesons: contain one  $b$  or  $\bar{b}$  [analogous to K's, D's]

$$B^+ = u\bar{b}$$

$$B^0 = d\bar{b}$$

$$B_s^0 = s\bar{b}$$

$$B_c^+ = c\bar{b} \text{ etc.}$$

Beautiful baryons

(1981)  $\Lambda_b = udb$

Top quark, long anticipated, not disc. until 1994 at Fermilab  
 Decays so quickly that no mesons are formed  
 $\Gamma \sim 1.4 \text{ GeV} \Rightarrow 5 \times 10^{-25} \text{ s} \ll 10^{-23} \text{ s}$

Why such a wide range of quark masses?

Why 3 generations?