


Massless particles

$$\boxed{p^2 = 0} \text{ for on-shell massless particles.}$$

[does that mean $p=0$?
of course not]

$$\text{ie } E^2 - |\vec{p}|^2 = 0$$

$$E = |\vec{p}| = \sqrt{p_x^2 + p_y^2 + p_z^2}$$



mass-shell : cone

[not hyperboloid]

$$\text{Since } \frac{\vec{p}}{E} = \vec{v}$$

massless particles obey $|\vec{v}| = 1$

ie must travel at the speed of light
(photons, gravitons)

For massless particle $T=E$ (all energy is kinetic)

($E = \gamma mc^2$, $\vec{p} = \gamma m \vec{v}$ are useless because $m \rightarrow 0$, $\gamma \rightarrow \infty$)

Photon energy & momentum depend, not on speed, but
on frequency & wavelength (wave number)

$$E = hf = \hbar \omega$$

$$|\vec{p}| = \frac{h}{\lambda} = \hbar k$$

$$E = |\vec{p}| \Rightarrow f = \frac{1}{\lambda} \text{ and } \omega = k$$

$$\text{ie } f = \frac{c}{\lambda} \text{ and } \omega = ck$$

- Physicists were skeptical of the photon concept
 until Arthur Compton did his expt on X-rays (1922)
 X-rays were known to be EM waves

Scattering of X-rays by electrons (~~Thomson~~ or Compton scattering)

Classical picture: electromagnetic waves of frequency f
 cause electrons to oscillate w/ frequency f .
 These accelerating electrons emit EM waves of frequency f .



Classical cross-section $\sigma = \frac{R}{F}$

R - rate at which energy is radiated by electron

F = flux of incident EM waves

From 1140, $F = \epsilon_0 c |\vec{E}|^2$

For 3120, Larmor formula $R = \frac{e^2 a^2}{6\pi \epsilon_0 c^3}$

$$\vec{a} = \frac{\vec{F}}{m_e} = \frac{e\vec{E}}{m_e} \Rightarrow R = \frac{e^4 |\vec{E}|^2}{6\pi \epsilon_0 m_e^2 c^3} \Rightarrow \sigma =$$

$$\Rightarrow \sigma = \frac{e^4}{6\pi \epsilon_0^2 (m_e c^2)^2} = \frac{8\pi}{3} \left(\frac{ke^2}{m_e c^2} \right)^2 = \frac{8\pi}{3} r_0^2$$

where $r_0 \equiv \frac{ke^2}{m_e c^2}$ = classical electron radius = $\frac{1.44 \text{ MeV fm}}{0.511 \text{ MeV}} = 2.8 \text{ fm}$

$$\sigma = 66.5 \text{ fm}^2 = \frac{2}{3} \text{ barn} = \text{Thomson cross-section}$$

[Thomson used it to measure # electrons in atoms]
 Nucleus too massive to radiate much.

I did this
 here in 2019
 but will
 probably do
 it later
 in QED
 instead
 not do this
 in future
 because not
 easy to do the
 quantum calc
 of Compton.

Compton, however, measured the wavelength of scattered X-rays and found it was slightly longer than incident X-rays

$$\lambda' > \lambda$$

We'll show that Compton shift $\Delta\lambda = \lambda' - \lambda$ is proportional to $\frac{h}{m_e c}$,


($\frac{h}{m_e c}$ called the Compton wavelength of the electron)

$$\frac{h}{m_e c} = \frac{2\pi\hbar c}{m_e c^2} = \frac{2\pi(197 \text{ MeV fm})}{0.511 \text{ MeV}} = 2400 \text{ fm} \approx 2.4 \times 10^{-12} \text{ m}$$

~~This is~~ an unnoticeably small shift for visible light $\lambda \sim 5 \times 10^{-7} \text{ m}$


[~~It is~~ is still small,] but measurable, for X-ray $\lambda \sim 10^{-10} \text{ m}$

Compton found
The shift depends on the scattering angle



$$\Delta\lambda = \lambda' - \lambda = \frac{h}{m_e c} (1 - \cos \theta)$$

Compton scattering can be understood as a collision
of a massless photon & a massive electron at rest
 $\gamma e^- \rightarrow \gamma e^-$



Photon gave some of its energy to the electron, so

$$E_{\gamma'} < E_\gamma$$

$$\text{Since } E = hf \Rightarrow f' < f$$

$$\text{Since } f = \frac{c}{\lambda} \Rightarrow \lambda' > \lambda$$

one can compute the Compton shift using
energy & momentum conservation

This is most elegantly done using 4-momentum



8-5

(Kinematics in fixed target frame)

$$p_\gamma = (E_\gamma, 0, 0, E_\gamma)$$

$$p_e = (m_e, 0, 0, 0)$$

$$p'_\gamma = (E'_\gamma, E'_\gamma \sin \theta, 0, E'_\gamma \cos \theta)$$

$$p'_e = (E'_e, -|\vec{p}'_e| \sin \phi, 0, |\vec{p}'_e| \cos \phi)$$

$$p_\gamma + p_e = p'_\gamma + p'_e$$

Rewrite this as

$$-p'_e + p_e = p'_\gamma - p_\gamma$$

Consider

$$\begin{aligned} (-p'_e + p_e)^2 &= p_e'^2 - 2p'_e \cdot p_e + p_e^2 \\ &= m_e^2 - 2(E'_e m_e - 0) + m_e^2 \\ &= 2m_e(m_e - E'_e) \stackrel{\substack{\uparrow \\ \text{energy cons.}}}{=} 2m_e(E_\gamma - E'_\gamma) \end{aligned}$$

Now consider

$$\begin{aligned} (p'_\gamma - p_\gamma)^2 &= p_\gamma'^2 - 2p'_\gamma \cdot p_\gamma + p_\gamma^2 \\ &= 0 - 2(E'_\gamma E_\gamma - E'_\gamma E_\gamma \cos \theta) \\ &= -2E_\gamma E'_\gamma (1 - \cos \theta) \end{aligned}$$

Equate them

$$-2m_e(E_\gamma - E'_\gamma) = -2E_\gamma E'_\gamma (1 - \cos \theta)$$

$$\frac{E_\gamma - E'_\gamma}{E_\gamma E'_\gamma} = \frac{1}{m_e} (1 - \cos \theta)$$

$$\frac{1}{E'_\gamma} - \frac{1}{E_\gamma} = \frac{1}{m_e c^2} (1 - \cos \theta)$$

↑
rest mass

$$E = hf = \frac{hc}{\lambda}$$

$$\Rightarrow \frac{\lambda'}{hc} - \frac{\lambda}{hc} = \frac{1}{m_e c^2} (1 - \cos \theta)$$

$$\lambda' - \lambda = \underbrace{\frac{h}{m_e c}}_{\text{Compton wavelength of an electron}} (1 - \cos \theta)$$

Compton wavelength of an electron


Max shift occurs γ backscattered X-rays ($\theta = \pi$)

$$(\Delta \lambda)_{\max} = \frac{2h}{m_e c} \approx 4.8 \times 10^{-12} \text{ m}$$

Compton used molybdenum K_α line $\Rightarrow 7 \times 10^{-11} \text{ m}$
 (17 keV)

(7% shift)

$$\gamma e^- \rightarrow e^-$$



$$(p_\gamma + p_e)^2 = p_e'^2$$

$$\underbrace{p_\gamma}_0^2 + 4p_\gamma p_e + \underbrace{p_e^2}_{m_e^2} = \underbrace{p_e'^2}_{m_e^2}$$


 $\Rightarrow$

$$p_\gamma p_e = 0$$

$$(E_\gamma m_e - 0) = 0$$


$$E_\gamma = 0$$

or lost in cm frame



violates conservation

Alternately



Can do this more formally using 4-vectors →

$$\begin{cases} E_\gamma + mc^2 = E_\gamma' + E_{e'} \\ p_\gamma = p_\gamma' \cos \theta + p_{e'} \cos \phi \\ 0 = p_\gamma' \sin \theta - p_{e'} \sin \phi \end{cases}$$

Eliminate ϕ :

$$(p_\gamma - p_\gamma' \cos \theta)^2 + (p_\gamma' \sin \theta)^2 = p_{e'}^2$$

$$p_\gamma^2 - 2p_\gamma p_\gamma' \cos \theta + p_\gamma'^2 = p_{e'}^2$$

$$\text{Energy} \Rightarrow (E_\gamma + mc^2 - E_\gamma')^2 = E_{e'}^2$$

$$E_\gamma^2 + E_\gamma'^2 + 2E_\gamma mc^2 - 2E_\gamma' mc^2 - 2E_\gamma E_\gamma' = E_{e'}^2 - (mc^2)^2$$

$$2E_\gamma mc^2 - 2E_\gamma' mc^2 - 2E_\gamma E_\gamma' (1 - \cos \theta) = 0 \quad \text{Subtract mm from energy}$$

$$\textcircled{a} \quad \frac{1}{E_\gamma'} - \frac{1}{E_\gamma} = \frac{1 - \cos \theta}{mc^2} \Rightarrow f(\theta) = \underline{\underline{1 - \cos \theta}}$$

$$\textcircled{b} \quad E = \frac{hc}{\lambda} \Rightarrow \lambda' - \lambda = \frac{h}{mc} (1 - \cos \theta)$$

$2.4 \times 10^{-12} \text{ m}$

$$\theta = \pi \Rightarrow \Delta \lambda = \underline{\underline{4.8 \times 10^{-12} \text{ m}}}$$

$$\lambda = \frac{hc}{17 \text{ KeV}} = \frac{1.24 \times 10^{-6} \text{ eV} \cdot \text{m}}{1.7 \times 10^4 \text{ eV}} = 7.3 \times 10^{-11} \text{ m}$$

$$\frac{\Delta \lambda}{\lambda} \sim 0.066$$

Alternate

Use 4-vectors to compute Compton scattering

$$p_e' = p_e + p_\gamma - p_\gamma'$$

$$(p_e')^2 = p_e^2 + p_\gamma^2 + p_\gamma'^2 + 2p_e \cdot p_\gamma - 2p_e \cdot p_\gamma' - 2p_\gamma \cdot p_\gamma'$$

$\uparrow \quad \uparrow \quad \quad \quad \uparrow \quad \uparrow$
 these cancel these vanish

so

$$p_e \cdot p_\gamma - p_e \cdot p_\gamma' = p_\gamma \cdot p_\gamma'$$


$$p_e = \begin{pmatrix} mc^2 \\ \vec{0} \end{pmatrix}, \quad p_\gamma = \begin{pmatrix} E \\ E\vec{n} \end{pmatrix}, \quad p_\gamma' = \begin{pmatrix} E' \\ E'\vec{n}' \end{pmatrix}$$

$$mc^2 E - mc^2 E' = EE' (1 - \underbrace{\vec{n} \cdot \vec{n}'}_{\cos \theta})$$

$$\frac{1}{E'} - \frac{1}{E} = \frac{1}{mc^2} (1 - \cos \theta)$$

Alternate:

$$p_e' - p_e = p_\gamma - p_\gamma'$$



$$2m^2 - 2p_e \cdot p_e' = -2p_\gamma \cdot p_\gamma'$$

$$2m^2 - 2mE_e = -2EE'(1 - \cos \theta)$$

$$\underbrace{E - E' = m}$$

$$-2m(E - E')$$

$$\frac{1}{E'} - \frac{1}{E} = \frac{1}{m} (1 - \cos \theta)$$

done in class